



Managing Immediate Implant Placement and Final Restoration Using Mucolock Concept of TTPHIL ALL TILT® Technique in Anterior Extraction Socket

Dr. Venkat Ratna Nag*, Dr. P Sarika¹, Dr. Smitha Daniel²

1. Dr P Sarika MDS (Pedodontia), Associate Professor, SB Patil Dental College, Bidar, India

Clinical Head, The Dental Specialists Clinic, Hyderabad, India.

2. Dr Smitha Daniel MDS (Prosthodontics), Prosthodontist, Institute for Dental Implantology, Hyderabad, India.

Corresponding Author: Dr. Venkat Ratna Nag, MDS(Prosthodontics), Director at Institute for Dental Implantology, Hyderabad, India. Professor, Department of Prosthodontics and Implantology, Panineeya Dental College, Hyderabad, India.

Copy Right: © 2021 Dr. Venkat Ratna Nag. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Received Date: July 06, 2021

Published date: July 15, 2021

Abstract

The placement of dental implants in the anterior region of the maxilla is challenging for clinicians because of the patient's high esthetic demands and difficult pre-existing anatomy. There are anatomic factors such as horizontal or vertical bone deficiencies and iatrogenic factors such as improper implant selection or dental implant malpositioning for an esthetic implant restoration. The immediate loading of implants is highly recommended especially in the anterior region where esthetic requirements are high.

In the literature, alternatives prove the effectiveness of immediate implant placement after tooth extraction that decreases the number of surgical procedures, reduces treatment time, and improved esthetics due to adequate osseointegration and much integration. This article presents a case with immediate implant placement and immediate loading using Tall Tilted Pin Hole Immediate Loading® (TTPHIL®) protocol in the anterior region with a fixed screw-retained prosthesis, satisfying the patient's functional and esthetic needs. TTPHIL® technique is considered as evidence-based dentistry, using tall and tilted implants with bicortical anchorage in a flapless, subcrestal manner delivering permanent prosthesis in 2-5 days. TTPHIL® concept is more advantageous in the esthetic zone as it provides soft tissue stability by creating a biologic seal.

Keywords: TTPHIL®, Tilted implant, Bicortical engagement, flapless, subcrestal, one time one abutment, mucolock.

Introduction

It is considered that placing implants after extraction in a single visit will reduce morbidity, treatment time, and treatment costs. This approach is called the "immediate implant and was first reported by Schulte et al in 1978.[1,2] Thereafter literature depicts several clinical reports with clinical and radiographic success.[3-11] The abundant experimental studies were performed on a light microscopic level in animals to confirm that osseointegration can be achieved following immediate implantation in immediate extraction sites.[12-14] In recent years, implant placement post-extraction in the esthetic zone for a single tooth is a frequent indication in implant therapy. [15]

The objective of implant treatment is an esthetic outcome, least number of surgical procedures, less pain and morbidity, low risk of complication, least treatment period, cost-effective to the patient.[16] TTPHIL® (Tall Tilted Pin Hole Immediate Loading®) is an advanced concept that has evolved from various philosophies in implantology (Basal, Pterygoid and Tilted) under immediate loading.[17-22] This technique utilizes the use of long tilted implants with bicortical engagement in the maxillary, nasal and pterygoid bone to maximize the success of rehabilitation. Tall implants (length: 16mm-25mm) and Tilted implants (30°-45°) with bicortical anchorage offer primary stability which supports osseointegration.[23] These implants bypass vital anatomical structures, reduce the cantilever effect, help in stress distribution and anterior-posterior spread.[24] The Pinhole (PH) placement of implant i.e flapless procedure reduces tissue trauma arising from flap reflection, maintaining soft tissue profile and reducing postoperative discomfort.[25,26] The implants are immediately loaded (IL) within 48 hours due to the good stability of implants attained from the bicortical anchorage.

Case Report



Figure 1: Preoperative frontal view



Figure 2: Drill passing through the surgical guide



Figure 3: Implant placed subcrestally




Figure 4: Abutment attached in the same appointment



Figure 5: Postoperative view of Zirconia crown restoration




Figure 6: Post operative follow up RVG at 2 years

A 35-year-old male patient presented with complained of pain in the front region for few days and also wanted replacement of missing back teeth of the upper jaw. The past medical history revealed no contraindication to dental treatment. Oral examination revealed acceptable oral hygiene, few missing teeth, pain on percussion in relation to 11(Figure 1). The periodontal evaluation revealed a thick periodontal biotype with generalized recession and probing depth of 3-4 mm. A thorough radiographic examination using orthopantomogram(OPG) and Cone Beam Computed Tomography (CBCT) was advised. The radiographic evaluation suggested a fractured # 11 with an unfavorable prognosis. The treatment plan was formulated and discussed with the patient, it was decided to extract tooth #11 followed by immediate implant placement and immediate loading using TTPHIL® technique.

Informed consent was obtained from the patient. Tooth extraction with immediate implant placement was planned. Under local anesthesia (2% lignocaine with 1:200000 adrenaline), atraumatic tooth extraction was performed. The extraction socket was thoroughly debrided and degranulated mechanically to remove all infected tissue followed by saline irrigation. Using a surgical guide, a single drill concept osteotomy was initiated through the extraction site with a pilot drill of diameter 1.2mm followed by a final single drill of 2.0 mm diameter (Figure 2). To obtain primary stability, osteotomies were performed at least 3 mm apical to the extraction socket. Bioline Surface Implant (3.75 mm diameter, 18 mm length) with an insertion torque of 45 N/cm² was placed subcrestally(Figure 3). The abutment was placed in the same appointment to avoid repeated detachments/attachments affecting the mucosal integration around it(Figure 4). Abutment level open impression was made. Shade selection was done and sent to a lab for zirconia crown fabrication. The bisque trial was done followed by shade confirmation and was sent to the lab for finishing the prosthesis. Using the CAD-CAM technique, the implant was loaded within 48 hours using a zirconia crown(Figure 5). The postoperative OPG was exposed showing implant and prosthesis. After 3 months, 1 year, and 2 years; the patient was recalled and it was noted to have exhibited good esthetic and stable bone levels(Figure 6). The tissues were healthy with no signs of inflammation.

Discussion

Whenever the dentists face a situation of tooth extraction in the esthetic zone, the decision is challenging in terms of esthetic success. Here, immediate extraction along with immediate implant placement is a highly predictable treatment plan. [27] The long treatment time and the use of a removable prosthesis during the treatment time may pose difficulty to many patients. Several authors considered 1-stage surgery, immediate implant placement, and immediate loading as the best option in case of replacement of anterior teeth.[28] When the atraumatic extraction is achieved with a flapless approach, the gingival architecture and blood supply to the buccal plate is preserved. In the literature, many techniques have been proposed to decrease the amount of hard and soft tissue collapse after extraction with different

ranges of successes. [29-31] Immediate implant placement and immediate loading using TTPHIL® technique has shown to maintain crestal bone levels through its concept of mucolock. This concept encompasses the advantages of flapless, subcrestal and one-time surgery to produce and maintain excellent esthetic results.[17] This provides a biologic seal around the abutment and maintains the hard and soft tissues. Nevertheless, its use in a larger number of cases and higher levels of evidence studies will provide further information and validity.

Conclusion

In this case report, the harmony of soft and hard tissue was achieved in the esthetically challenging situation by immediate implant placement and immediate loading using TTPHIL® technique. Minimally invasive surgical procedures, lesser treatment time with minimum post-extraction complications and preservation of gingival aesthetics are a boon to the patient. However proper case selection, diagnosis and treatment planning, meticulous post-operating care are very essential long term success. Thus, the result of this case report has the potential to be recommended for similar cases.

References

1. Wilson T, Weber H. "Classification of and therapy for areas of deficient bony implant placement". Int J Periodont Rest Dent 1993;13:451-459.
2. Schulte W. Kleineikenscheidt H, Linder K. Schareyka R. "The Tübingen immediate implant in clinical studies". DtschZahnärztl Z. 1978;5:348-359.
3. Werbitt M, Goldberg P. "The immediate implant: Bone preservation and bone regeneration In Periodont Rest" Dent 1992;12:206-217.
4. Becker W, Dahlin C, Becker BE, Lekholm U, Van Steenberghe D. Higuchi-Kultje C. "The use of PTFE barrier membranes for bone promotion around titanium implants placed into extraction socket retrospective multicenter study". Int J Oral Maxillofac Implants 1994;9:31-40.
5. Nyman S. Lang NP, Buser D, Brügger U. "Bone regeneration adjacent to titanium dental implants guided tissue regeneration: A report of two cases". Int J Oral Maxillofac Implants 1990;5:9-14.
6. Lazzara RJ. "Immediate implant placement into extraction sites: Surgical and restorative advantage" J Periodont Rest Dent 1989;9:333-343.
7. Gelb DA. "Immediate implant surgery: 3 year retrospective evaluation of 50 consecutive cases". Int Maxillofac Implants 1993;8:388-399.

8. Watzek G, Haider R, Mensdorff-Pouilly N, Haas R. "Immediate implants and delayed implants: complete restoration of the jaw following extraction of all residual teeth: A retrospective comparing different types of serial implantation". *Int J Oral Maxillofac Implants* 1995;10:561-56
9. Rosenquist B, Grenthe B. "Immediate placement of implants into extraction sockets: Report a procedure". *Int J Oral Maxillofac Implants* 1991;6:277-284.
10. Langer B, Langer L., Herrmann I, Jorneus L. "The wide fixture: A solution for special bone situation a rescue for the compromised implant". Part 1. *Int J Oral Maxillofac Implants* 1993;8:400-408.
11. Lang N, Bragger U, Hammerle C, Sutter F. "Immediate transmucosal implants using the principle of guided tissue regeneration. 1. rationale, clinical procedures and 30-month results". *Clin Oral Implants Res* 1994;5:154-163.
12. Lundgren D, Ryland H, Andersson M, Johansson C, Albrektsson T. "Healing-in of root analog implants placed in extraction sockets: An experimental study in beagle dogs". *Clin Oral Surg Res* 1992;3:136-144.
13. Knox R, Caudell R, Meffert R. "Histologic evaluation of dental endosseous implants placed in surgically created extraction defects". *Int J Periodont Rest Dent* 1991;11:365-376.
14. Wilson TG, Schenk R, Med, Buser D et al. "Implant placed in immediate extraction sites: A report of histologic and histometric analyses of human biopsies". *Int J of Oral Maxillofac Implant* 1998;13(3):3
15. Brugger OE, Bornstein MM, Kuchler U, Janner SF, Chappuis V, Buser D. "Implant therapy in a surgical specialty clinic: an analysis of patients, indications, surgical procedures, risk factors, and early failures" *Int J Oral Maxillofac Implants* 2015; 30: 151-160.
16. Buser D, Chen ST. Implant placement in postextraction sites. In: Buser D. editor. 20 years of guided bone regeneration in implant dentistry, 2nd edition. Chicago: Quintessence Publishing Co., Inc. 2009. 153-194.
17. Nag PVR, Dhara V. et al. "Treatment of the Complete Edentulous Atrophic Maxilla: The Tall Tilted Pin Hole Placement Immediate Loading (TTPHIL)-ALL TILT™ Implant Option". *J Contemp Dent Pract* 2019;20(6):754-763.
18. VenkatRatna Nag P. "Immediate Implant Placement and Loading with Tall and Tilted Pinhole Immediate Loading (TTPHIL) Technique". *Guident Sep* 2017:26-27
19. Venkat Nag P, Sarika P, Pavankumar, A. (2017). "TTPHIL-ALL TILT™ Concept - An Innovative in Immediate Functional Loading Implant Placement in Maxilla". *Scholars Journal of Dental Technique in Sciences*. 4(9),397-399.

20. Venkat Nag P, Sarika P. Khan R, Bhagwatkar T. (2018). "Tall and tilted pin hole immediately loaded implants (TTPHIL) technique for maxillary arch rehabilitation". International Journal of Research and Review, 5:104-110.
21. Venkat Nag P, Sarika P, Khan R. Bhagwatkar T. (2018). "Immediate implantation and loading in just two days with TIPHIL technique using CAD/CAM Prosthesis". International Journal of Applied Dental Sciences, 4(3): 209-213.
22. VenkatRatna Nag. Sarika P.Bhagwatkar T.Dhara V. "Tilted Implant Placement for Rehabilitation of Severe Atrophic Maxilla Supported by Finite Element Analysis". The Journal of Implant & Advanced Clinical Dentistry 2019;11(1):14-26.
23. Andrea H, Wook-Jin S, Ryan W. "Comparison of initial implant stability of implants placed using bicortical fixation, Indirect sinus elevation, and unicortical fixation". Int J of Oral Max Implants 2016.31:2:459-468)
24. Leonard K, Mikael K, Rangert B. "Tilting of Posterior mandibular and maxillary implants for improved prosthesis support". Int J of Oral Max Implants 2000: 405-414
25. Brodala N. "Flapless Surgery and Its Effect on Dental Implant Outcomes". Int J Oral Maxillofac Implants 2009; 24:118-125.
26. Laverty DP, Buglass J, Patel A. "Flapless Dental Implant Surgery and Use Of Cone Beam Computer Tomography Guided Surgery". British Dental Journal 2018;224:601-611.
27. Wael G. "A combination of buccal plate preservation and dual zone therapeutic technique in immediate implants: a one year follow up". J Oral Implantology 5:42:484-89.
28. Park JB. "Immediate placement of dental implants into fresh extraction socket in the maxillary anterior region. A case report". J of Oral Implantology 2010;36(2):153-57.
29. Cosyn J, Eghbali A, De Bruyn H, Collys K, Cleymaet R, De Rouck T. "Immediate single-tooth implants in the anterior maxilla: 3-year results of a case series on hard and soft tissue response and aesthetics". J ClinPeriodontol. 2011;38:746-753.
30. Tsuda H, Rungcharassaeng K, Kan JY, Roe P, Lozada JL, Zimmerman G. "Peri-implant tissue response following connective tissue and bone grafting in conjunction with immediate single-tooth replacement in the esthetic zone: a case series". Int J Oral Maxillofac Implants. 2011;26:427-436.
31. Migliorati M. Amorfini L, Signori A, Biavati AS, Benedicenti S. "Clinical and aesthetic outcome with post-extractive implants with or without soft tissue augmentation: a 2-year randomized clinical trial". Clin Implant Dent Relat Res. 2015;17:983-995.